(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 1 September 2005 (01.09.2005)

(10) International Publication Number WO 2005/080428 A2

(51) International Patent Classification7:

C07K 16/00

(21) International Application Number:

PCT/EP2005/001307

(22) International Filing Date: 9 February 2005 (09.02.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10/778,915

13 February 2004 (13.02.2004) US

- (71) Applicant (for all designated States except US): MI-CROMET AG [DE/DE]; Staffelseestrasse 2, 81477 München (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): PETERS, Malte [DE/DE]; c/o Micromet AG, Staffelseestrasse 2, 81477 München (DE). LOCHER, Mathias [DE/DE]; c/o Micromet AG. Staffelseestrasse 2, 81477 München (DE). PRANG, Nadja (DE/DE); c/o Micromet AG, Staffelseestrasse 2, 81477 München (DE). QUADT, Cornella (DE/DE); c/o Micromet AG, Staffelseestrasse 2, 81477 München (DE).

 8 Micromet AG, Staffelseestrasse 2, 81477 München

- (74) Agent: DEHMEL, Albrecht; Dehmel & Bettenhausen, Herzogspitalstr. 11, 80331 München (DE).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU. LV, MA, MD, MG, MK, MN. MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH. PL. PT. RO. RU. SC. SD. SE. SG. SK. SL. SY. TJ. TM. TN. TR. TT. TZ. UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

without international search report and to be republished upon receipt of that report

the patient a human immunoglobulin specifically binding to the human EpCAM antigen, the immunoglobulin exhibiting a serum half-life of at least 15 days, the method comprising the step of administering the immunoglobulin no more frequently than once every week, preferably no more frequently than once every two weeks.